



## Andro Balance References

- Sakawa K, et al. *Effects of certain pollen-extract (Cernilton) on inflammatory cytokines in sex-hormone-induced nonbacterial prostatitis rats.* Hinyokika Kyo 2001 Jul;47(7):459-465.
- Baker AM, et al. *Expression of antioxidant enzymes in human prostatic adenocarcinoma.* Prostate 1997;32(4):229-233.
- Bartlet A, et al. *Efficacy of Pygeum africanum extract in the medical therapy of urination disorders due to benign prostatic hyperplasia:evaluation of objective and subjective parameters. A placebo-controlled double-blind multicenter study.* Wein Klin Wochenschr 1990;23;102(22):667-673.
- Bartsch G, et al. *Dihydrotestosterone and the concept of 5alpha-reductase inhibition in human benign prostatic hyperplasia.* Eur Urol. 2000;37(4):367-80.
- Becker H, Eveling L. *Conservative treatment of benign prostatic hyperplasia (BPH) with Cernilton N - Results of a placebo-controlled double-blind study.* In: Benign Prostatic Diseases, Vahlensiek W, Rutishauser G, Eds. Thieme Medical Publishersm New York, pp:1166-1174.
- Berggren MM, et al. *Effect of selenium on rat thioredoxin reductase activity: increase by supranutritional selenium and decrease by selenium deficiency.* Biochem Pharmacol 1999 Jan 15;57(2):187-193.
- Bracher F. *Phytotherapy of benign prostatic hyperplasia.* Urologe A 1997;36(1):10-7.
- Buck AC, et al. *Treatment of outflow tract obstruction due to benign prostatic hyperplasia with the pollen extract, cernilton. A double-blind, placebo-controlled study.* Br J Urol 1990;66(4):398-404.
- Carbin BE, et al. *Treatment of benign prostatic hyperplasia with phytosterols.* Br J Urol 1990;66(6):639-41.
- Clark LC, et al. *Decreased incidence of prostate cancer with selenium supplementation: results of a double-blind cancer prevention trial.* Br J Urol 1998 May;81(5):730-734.
- Clinton SK. *Lycopene: chemistry, biology, and implications for human health and disease.* Nutr Rev 1998;56(2 Pt 1):35-51.
- Gerster H. *The potential role of lycopene for human health.* J Am Coll Nutr 1997;16(2):109-26.
- Habib FK, et al. *Identification of a prostate inhibitory substance in a pollen extract.* Prostate 1995;26(3):133-139.
- Horton R. *Benign prostatic hyperplasia. A disorder of androgen metabolism in the male.* J Am Geri Soc 1984;32:380-385.
- Hryb DJ, et al. *The effect of extracts of the roots of the stinging nettle (Urtica dioica) on the interaction of SHBG with its receptor on human prostatic membranes.* Planta Med 1995 Feb;61(1):31-2.

- Ishani A, et al. *Pygeum africanum* for the treatment of patients with benign prostatic hyperplasia: a systematic review and quantitative meta-analysis. *Am J Med* 2000;109(8):654-664.
- Judd AM, et al. Zinc acutely, selectively and reversibly inhibits pituitary prolactin secretion, *Brain Res* 1984;294(1):190-192.
- Konrad L, et al. Antiproliferative effect on human prostate cancer cells by a stinging nettle root (*Urtica dioica*) extract. *Planta Med* 2000;66(1):44-7.
- Krzeski T, et al. Combined extracts of *Urtica dioica* and *Pygeum africanum* in the treatment of benign prostatic hyperplasia: double-blind comparison of two doses. *Clin Ther* 1993;15(6):1011-1020.
- Leake A, et al. The effect of zinc on the 5 alpha-reduction of testosterone by the hyperplastic human prostate gland. *J Steroid Biochem* 1984;20(2):651-655.
- Mathe G, et al. A *Pygeum africanum* extract with so-called phyto-estrogenic action markedly reduces the volume of true and large prostatic hypertrophy. *Biomed Pharmacother* 1995;49(7-8):341-3.
- Menter DG, et al. Selenium effects on prostate cell growth. *Cancer Epidemiol Biomarkers Prev* 2000;9(11):1171-1182.
- Paubert-Braquet M, et al. Effect of *Pygeum africanum* extract on A23187-stimulated production of lipoxygenase metabolites from human polymorphonuclear cells. *J Lipid Mediat Cell Signal* 1994;9(3):285-290.
- Picurelli L, et al. Determination of Fe, Mg, Cu, and Zn in normal and pathological prostatic tissue. *Actas Urol Esp* 1991 Jul;15(4):344-350.
- Rao AV, Agarwal S. Bioavailability and in vivo antioxidant properties of lycopene from tomato products and their possible role in the prevention of cancer. *Nutr Cancer* 1998;31(3):199-203.
- Rao AV, et al. Serum and tissue lycopene and biomarkers of oxidation in prostate cancer patients: a case-control study. *Nutr Cancer* 1999;33(2):159-64.
- Rugendorff EW, et al. Results of treatment with pollen extract (cernilto N) in prostatodynia and chronic prostatitis. In: *Benign Prostatic Diseases*, Vahlensiek W, Rutishauser G, Eds. Thieme Medical Publishers, New York, pp:73-83.
- Schiebel-Schlosser G, Friederich M. Phytotherapy of BPH with pumpkin seeds - A multicentric clinical trial. *Zeit fur Phytother* 1998;19:71-76.
- Tunn S, Kreig M. Alterations in the intraprostatic hormonal metabolism by the pollen extract Cernilton. In: *Benign Prostatic Diseases*, Vahlensiek W, Rutishauser G, Eds. Thieme Medical Publishers, New York, pp:109-114.
- Wallace AM, Grant JK. Effect of zinc on androgen metabolism in the human hyperplastic prostate. *Biochem Soc Trans* 1975;3(4):540-542.
- Yasumoto R, et al. Clinical evaluation of long-term treatment using cernitin pollen extract in patients with benign prostatic hyperplasia. *Clin Ther* 1995;17(1):82-87.